# antibodies -online.com







# anti-FRAT2 antibody (AA 131-230) (Alexa Fluor 750)



$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
	· · · · · · · · · · · · · · · · · · ·	
Target:	FRAT2	
Binding Specificity:	AA 131-230	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FRAT2 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FRAT2	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	FRAT2
Alternative Name:	FRAT2 (FRAT2 Products)

## **Target Details**

Synonyms: Frequently rearranged in advanced T-cell lymphomas 2 antibodyGSK-3 binding		
protein FRAT2, FRAT2_HUMAN.		
Background: FRAT1 and FRAT2 were originally characterized as proteins frequently rearranged		
in advanced T cell lymphoma, and they have since been identified as proto-oncogenes involved		
in tumorigenesis. These proteins share significant homology with the Xenopus glycogen		
synthase kinase-3 (xGSK-3) binding protein, which is designated GBP and is essential for the		
formation of the dorsal-ventral axis during embryonic development. Establishment of these		
embryonic axes is mediated by the Wnt intracellular signaling pathway. Wnt signaling is		
regulated in part by the activity of GSK-3, which phosphorylates and thereby facilitates the		
degradation of ?catenin. GBP binds to GSK-3 and inhibits this phosphorylation, resulting in the		
accumulation of ?catenin and the subsequent transcription of Wnt target genes. Like GBP,		
FRAT2 has been shown to bind xGSK-3, suggesting that FRAT1 and FRAT2 may be GSK-3		
regulatory proteins.		
23401		
IF(IHC-P) 1:50-200		
IF(IHC-F) 1:50-200		
IF(ICC) 1:50-200		
For Research Use only		
Liquid		
1 μg/μL		
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
50 % Glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
handled by trained staff only.		
-20 °C		
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
12 months		