# antibodies .- online.com





# anti-FUT6 antibody (AA 101-200) (AbBy Fluor® 750)



( )	1 /	$\sim$	rv	11/	11	Α
	1//	⊢	I \/	16	٦,	/\

Quantity:	100 μL	
Target:	FUT6	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FUT6 antibody is conjugated to AbBy Fluor® 750	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FUT6/FucT-VI	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	FUT6	
Alternative Name:	FUT6 (FUT6 Products)	
Background:	kground: Synonyms: Alpha 1,3 fucosyltransferase, EC=2.4.1.65, FCT3A, Fuc TVI, Fucosyltransferase	

Fucosyltransferase VI, FucT VI, Galactoside 3 L fucosyltransferase, FUT6\_HUMAN.

Background: Glycosyltransferases that mediate the regio- and stereoselective transfer of sugars, such as the fucosyltransferases, determine cell surface-carbohydrate profiles, which is an essential interface for biological recognition processes. Fucosyltransferases catalyze the covalent association of fucose to different positional linkages in sugar acceptor molecules. The carbohydrate moieties generated and covalently attached to cell surfaces are necessary to ensure a surface contour that satisfies physiological roles, which are reliant on adhesion molecules such as Selectins (1-3). Hematopoietic lineages rely on Fucosyltransferases to confer a surface carbohydrate phenotype, which mediates proper cell adhesion molecule recruitment and cell trafficking (4-6).

Gene ID:

2528

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months