

## Datasheet for ABIN5003143

## anti-GALM antibody (AA 251-342) (AbBy Fluor® 680)



Go to Product page

Overview	
Quantity:	100 μL
Target:	GALM
Binding Specificity:	AA 251-342
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GALM antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human GALM/Galactose mutarotase
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	GALM
Alternative Name:	GALM/Galactose mutarotase (GALM Products)

## **Target Details**

Expiry Date:

Synonyms: Aldose 1 epimerase, Aldose 1-epimerase, BLOCK25, Galactomutarotase, Galactose
mutarotase, Galm, GALM_HUMAN, IBD1.
Background: Galactose Mutarotase is a member of the aldose epimerase family and is involved
in hexose metabolism. Through its catalytic activity, Galactose Mutarotase converts beta-
aldose to alpha-aldose on several sugars, including D-glucose, L-arabinose and D-xylose. Found
in the cytoplasm of most cells, Galactose Mutarotase plays a key role in galactose metabolism
by catalyzing the conversion of beta-D-galactose to alpha-D-galactose. The enzyme contains
two residues, Glu 304 and His 170, that are critical for catalysis, as well as His 96 and Asp 243,
which are important for proper substrate recognition by the active site. No known diseases
have been associated with mutations in the Galactose Mutarotase gene, although inhibition of
Galactose Mutarotase activity could potentially be associated with a build-up of unmetabolized
sugars during metabolism.
130589
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months