

Datasheet for ABIN5003251

anti-GBA3 antibody (AA 151-250) (AbBy Fluor® 680)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GBA3
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GBA3 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GBA3/CBG
Isotype:	IgG
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	GBA3
Alternative Name:	GBA3/CBG (GBA3 Products)
Background:	Synonyms: CBG, CBGL1, Cytosolic beta glucosidase, Cytosolic beta glucosidase like protein 1,

Target Details

Cytosolic beta-glucosidase, Cytosolic beta-glucosidase-like protein 1, GBA3, GBA3_HUMAN, GLUC, Glucosidase beta acid 3, Klotho related protein.

Background: CBG is a monomeric enzyme involved in the absorption and metabolism of flavonoid glucosides. CBG is found predominately in the liver, but is also located in tissues such as spleen, small intestine and kidney. Through its catalytic activity, CBG is able to hydrolyze a variety of glycosides including phytoestrogens, cyanogens, and flavonols. Although its catalytic activity extends to many dietary flavonoids, CBG has increased specificity for hydrophobic aglycones such as beta-D-glucoside and beta-D-galactoside. Hydrolysis is inhibited by sodium taurocholate and glucosyl-sphingosine, both of which regulate CBG enzymatic activity.

Deficiencies in CBG have been implicated in Gaucher's disease, a lysosomal storage disease that causes a build up of fatty material in the spleen, liver, lung and kidneys.

Gene ID: 57733

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months