

Datasheet for ABIN5003374

anti-SMN1 antibody (AA 31-100) (AbBy Fluor® 680)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	SMN1
Binding Specificity:	AA 31-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMN1 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Gemin 1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	SMN1
Alternative Name:	Gemin 1/SMA (SMN1 Products)

Target Details

Background:	Synonyms: Component of gems 1, Gemin 1, Gemin-1, Gemin1, SMA 1, SMA 2, SMA 3, SMA 4, SMA, SMA1, SMA2, SMA3, SMA4, SMN 1, SMN, SMN-1, SMN_HUMAN, SMN1, SMN2, SMNT, Survival motor neuron protein, Survival of motor neuron 1 telomeric, survival of motor neuron 1, Survival of motor neuron 1, telomeric, T-BCD541, BCD541, SMN_HUMAN. Background: Spinal muscular atrophy (SMA) is an autosomal recessive neurodegenerative disease characterized by loss of motor neurons in the spinal cord. SMA is caused by deletion or loss-of-function mutations of SMN (survival of motor neuron) gene. SMN, also known as Gemin1, SMN1, SMNT and BCD541, exists as four isoforms produced by alternative splicing. SMN is oligomeric and forms a complex with Gemin2 (formerly SIP1), Gemin3 (a DEAD box RNA helicase), Gemin4, Gemin5 and Gemin6, as well as several spliceosomal snRNP proteins. The SMN complex plays an essential role in spliceosomal snRNP assembly in the cytoplasm and is required for pre-mRNA splicing of the nucleus. The SMN complex is found in both the cytoplasm and the nucleus. The nuclear form is concentrated in subnuclear bodies called gems (gemini of the coiled bodies). Cytoplasmic SMN interacts with spliceosomal Sm proteins and facilitates their assembly onto U snRNAs, and nuclear SMN mediates recycling of pre-mRNA splicing factors. Nearly identical telomeric and centromeric forms of SMN encode the same protein, however, only mutations in the telomeric form are associated with the disease-state SMA. SMN is expressed in a wide variety of tissues including brain, kidney, liver, spinal cord and moderately in skeletal and cardiac muscle.
-------------	--

Pathways:	Ribonucleoprotein Complex Subunit Organization
-----------	--

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months