antibodies -online.com





anti-GFM1 antibody (AA 401-500) (Alexa Fluor 750)



_		
()Ver	view	

0	
Quantity:	100 μL
Target:	GFM1
Binding Specificity:	AA 401-500
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GFM1 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GFM1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	GFM1
Alternative Name:	GFM1 (GFM1 Products)

Target Details

Background:	Synonyms: COXPD1, EF-Gmt, EFG, EFG1, EFGM, EFGM_HUMAN, EGF1, Elongation factor G 1,
-	Elongation factor G 1 mitochondrial, Elongation factor G, Elongation factor G1, FLJ12662,
	FLJ13632, FLJ20773, G elongation factor mitochondrial 1, G translation elongation factor
	mitochondrial, GFM 1, GFM, gfm1, hEFG1, mEF G 1, mEF-G 1, mEFG 1, mitochondrial,
	Mitochondrial elongation factor G1.
	Background: Mitochondrial GTPase that catalyzes the GTP-dependent ribosomal translocation
	step during translation elongation. During this step, the ribosome changes from the pre-
	translocational (PRE) to the post-translocational (POST) state as the newly formed A-site-bound
	peptidyl-tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively.
	Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and
	conformational changes in the ribosome. Does not mediate the disassembly of ribosomes from
	messenger RNA at the termination of mitochondrial protein biosynthesis.
Gene ID:	85476
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Otomorphi	
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months