

Datasheet for ABIN5003602

**anti-PRKCSH antibody (AA 101-200) (AbBy Fluor® 680)**[Go to Product page](#)

## Overview

|                      |                                                                                                                                |
|----------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Quantity:            | 100 µL                                                                                                                         |
| Target:              | PRKCSH                                                                                                                         |
| Binding Specificity: | AA 101-200                                                                                                                     |
| Reactivity:          | Human                                                                                                                          |
| Host:                | Rabbit                                                                                                                         |
| Clonality:           | Polyclonal                                                                                                                     |
| Conjugate:           | This PRKCSH antibody is conjugated to AbBy Fluor® 680                                                                          |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |                                                                                |
|-----------------------|--------------------------------------------------------------------------------|
| Immunogen:            | KLH conjugated synthetic peptide derived from human Glucosidase 2 subunit beta |
| Isotype:              | IgG                                                                            |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Pig, Horse                                             |
| Purification:         | Purified by Protein A.                                                         |

## Target Details

|                   |                                                                              |
|-------------------|------------------------------------------------------------------------------|
| Target:           | PRKCSH                                                                       |
| Alternative Name: | Glucosidase 2 subunit beta ( <a href="#">PRKCSH Products</a> )               |
| Background:       | Synonyms: 80K-H protein, AGE-binding receptor 2, AGE-R2, G19P1, GLU2B_HUMAN, |

## Target Details

Glucosidase 2 subunit beta, Glucosidase II beta subunit, Glucosidase II subunit beta, Hepatocystin, PCLD, PKCSH, PLD1, PRKCSH, Protein kinase C substrate 60.1 kDa protein heavy chain, Protein kinase C substrate 80 Kda protein, Protein kinase C substrate 80K-H, Protein kinase C substrate, 80 Kda protein.

Background: Trimming of glucoses from N-linked core glycans on newly synthesized glycoproteins occurs sequentially through the action of Glucosidases I and II in the endoplasmic reticulum (ER). Glucosidase II is an ER-localized enzyme that contains a and b subunits (Glucosidase IIa and Glucosidase IIb) which form a defined heterodimeric complex. Glucosidase IIa is the catalytic core of the enzyme and can function independently of the b subunit. The sequence of Glucosidase IIb encodes protein rich in glutamic and aspartic acid with a putative ER retention signal (HDEL) at the C-terminus. The phosphorylated form of Glucosidase IIb is localized in the plasma membrane and is highly expressed in FGF-stimulated fibroblasts and epidermal carcinoma cells. Glucosidase IIb was first purified from a human carcinoma cell line as a potential substrate for protein kinase C. Through the HDEL signal at the C-terminus, Glucosidase IIb retains the complete complex in the ER.

Gene ID: 5589

Pathways: [Cellular Glucan Metabolic Process](#), [Methionine Biosynthetic Process](#)

## Application Details

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

---

|                  |                                                                                   |
|------------------|-----------------------------------------------------------------------------------|
| Storage:         | -20 °C                                                                            |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date:     | 12 months                                                                         |