antibodies .- online.com





anti-Golgin A3 antibody (AA 951-1050) (Alexa Fluor 750)



Go to Product page

\sim					
	1/6	⊃r	\/I	\triangle	٨/

Quantity:	100 μL	
Target:	Golgin A3 (GOLGA3)	
Binding Specificity:	AA 951-1050	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Golgin A3 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Golgin 160
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Golgin A3 (GOLGA3)
Alternative Name:	Golgin 160 (GOLGA3 Products)
Background:	Synonyms: GCP170, GOLGA 3, GOLGA3, Golgi autoantigen, Golgi complex associated protein of

170 kDa, Golgi complex-associated protein of 170 kD, Golgi membrane associated protein, Golgi peripheral membrane protein, Golgin 160, Golgin subfamily a 3, HGNC:4426, Male enhanced antigen 2, MEA2, SY2/SY10 protein, GOGA3_HUMAN.

Background: The Golgi apparatus consists of a series of stacked, flattened membrane stacks called cisternae that are involved in the transport of lipids and proteins in the secretory pathway and are important for Golgi-microtubule interaction. Golgin 160 is a 1,498 amino acid protein that localizes to both the cytoplasm and to the Golgi apparatus and contains a series of coiled-coil domains. Expressed in a variety of tissues, including heart, liver, testis, kidney, lung and salivary gland, golgin 160 functions as a homodimer that interacts with GOLGA7 and is thought to be involved in maintaining Golgi structure and may play a role in nuclear transport and Golgi apparatus localization. Multiple isoforms of golgin 160 exist due to alternative splicing events.

Gene ID:

2802

Pathways:

SARS-CoV-2 Protein Interactome

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months