antibodies.com

## Datasheet for ABIN5003816 anti-GPBP1L1 antibody (AA 261-360) (Alexa Fluor 750)



Overview

Quantity:	100 µL
Target:	GPBP1L1
Binding Specificity:	AA 261-360
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPBP1L1 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GPBP1L1
lsotype:	lgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

Target:	GPBP1L1
Alternative Name:	GPBP1L1 (GPBP1L1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5003816 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Background:       Synonyms: GC-rich promoter-binding protein 1-like 1, GPBL1_HUMAN, GPBP1L1, SF         Vasculin-like protein 1.       Background: Transcription factors are required for the initiation of transcription. The transcription by binding to DNA at specific nucleotide sequences within promoters enhancers. Transcription factors, which may also bind to RNA polymerase or to oth transcription factors, are involved in the preinitiation complex formation. Upstream transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducibl require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.         Gene ID:       60313         Application Notes:       IF(IHC-P) 1:50-200         IF(IHC-F) 1:50-200       IF(IHC-F) 1:50-200	Y
Background: Transcription factors are required for the initiation of transcription. The transcription by binding to DNA at specific nucleotide sequences within promoters enhancers. Transcription factors, which may also bind to RNA polymerase or to oth transcription factors, are involved in the preinitiation complex formation. Upstream transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	<sup>&gt;</sup> 192,
transcription by binding to DNA at specific nucleotide sequences within promoters enhancers. Transcription factors, which may also bind to RNA polymerase or to oth transcription factors, are involved in the preinitiation complex formation. Upstream transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	
enhancers. Transcription factors, which may also bind to RNA polymerase or to oth transcription factors, are involved in the preinitiation complex formation. Upstream transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	ey regulate
transcription factors, are involved in the preinitiation complex formation. Upstream transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	and
transcription factors and inducible transcription factors bind upstream of the initiat repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	ier
repress or stimulate transcription. Upstream factors are unregulated, while inducible require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	
require inhibition or activation. GPBP1L1 (GC-rich promoter binding protein 1-like 1) as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	ion site to
as vasculin-like protein 1 or SP192, is a 474 amino acid protein belonging to the vas Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.Gene ID:60313Application DetailsIF(IHC-P) 1:50-200	e factors
Localizing to nucleus, GPBP1L1 may function as a transcription factor. The gene er         GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.         Gene ID:       60313         Application Details         Application Notes:       IF(IHC-P) 1:50-200	, also known
GPBP1L1 maps to human chromosome 1p34.1 and mouse chromosome 4 D1.         Gene ID:       60313         Application Details         Application Notes:       IF(IHC-P) 1:50-200	sculin family.
Gene ID:     60313       Application Details     IF(IHC-P) 1:50-200	ncoding
Application Details Application Notes: IF(IHC-P) 1:50-200	
Application Notes: IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
Restrictions: For Research Use only	
Handling	
Format: Liquid	
Concentration: 1 µg/µL	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pro	oclin300 and
50 % Glycerol.	
Preservative: ProClin	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which	h should be
handled by trained staff only.	
Storage: -20 °C	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date: 12 months	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5003816 | 03/08/2024 | Copyright antibodies-online. All rights reserved.