antibodies -online.com





anti-GPLD1 antibody (Alexa Fluor 750)



Go to Product page

\sim	
()\/\	rview
\circ	1 410 44

Quantity:	100 μL
Target:	GPLD1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPLD1 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GPLD1/GPI-PLD
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	GPLD1
Alternative Name:	GPLD1 (GPLD1 Products)
Background:	Synonyms: Glycoprotein phospholipase D, Glycosyl-phosphatidylinositol-specic phospholipase D, GPI-PLD, GPI-specic phospholipase D, GPIPLD, GPIPLDM, GPLD1, PHLD_HUMAN,
	Phosphatidylinositol glycan specic phospholipase D, Phosphatidylinositol-glycan-specic
	phospholipase D, PI-G PLD, PIGPLD, PIGPLD1.

Background: Phosphatidylinositol-glycan-specific phospholipase D (GPI-PLD) is a high-density lipoprotein-associated protein found on chromosome 6p22 that specifically hydrolyzes the inositol phosphate linkage in proteins anchored by phosphatidylinositol-glycans (PI-Gs). GPI-PLD is found in serum, liver, cerebrospinal fluid and in milk. The majority of plasma GPI-PLD appears to be specifically associated with a small, discrete and minor fraction of lipoproteins containing apoA-I and apoA-IV. Serum GPI-PLD activity is reduced over 75 % in systemic inflammatory response syndrome and the downregulation of GPI-PLD could play an important role in the control of proinflammatory responses.

Gene ID:

2822

Pathways:

Inositol Metabolic Process, Regulation of Carbohydrate Metabolic Process

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months