
# antibodies .- online.com





## anti-HCN3 antibody (Alexa Fluor 680)



Go to Product page

| ( ) | 11/0   | K\ /   | iew         |
|-----|--------|--------|-------------|
|     | $\cup$ | 'I V/I | $I \cap VV$ |
|     |        |        |             |

| Quantity:    | 100 μL  |  |
|--------------|---|--|
| Target:      | HCN3  |  |
| Reactivity:  | Human, Mouse, Rat   |  |
| Host:        | Rabbit  |  |
| Clonality:   | Polyclonal  |  |
| Conjugate:   | This HCN3 antibody is conjugated to Alexa Fluor 680                             |  |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |
|              |   |  |

#### **Product Details**

| Immunogen:        | KLH conjugated synthetic peptide derived from human HCN3 |  |
|-------------------|--|--|
| Isotype:          | IgG  |  |
| Cross-Reactivity: | Human, Mouse, Rat  |  |
| Purification:     | Purified by Protein A.                                   |  |

## Target Details

| Target:           | HCN3   |
|-------------------|--|
| Alternative Name: | HCN3 (HCN3 Products)   |
| Background:       | Synonyms: Hcn3, HCN3_HUMAN, hyperpolarization activated cyclic nucleotide-gated potassium channel 3, KIAA1535, potassium/sodium hyperpolarization-activated cyclic |
|                   | nucleotide-gated channel 3.  |
|                   | Background: Hyperpolarization-activated, cyclic nucleotide-binding channels (HCN) are voltage-   |

#### **Target Details**

gated cation channels that are activated by direct binding of intracellular cyclic nucleotides. The HCN family consists of four members (HCN1-4), each with a core transmembrane segment domain and a C-terminal 120 amino-acid cyclic nucleotide-binding domain motif. HCN channels are expressed in the brain, heart, thalamus and testis. The pacemaker properties of HCN channels contribute to spontaneous rhythmic activity in the brain and heart. HCN3 contains a segment characterized by a series of positively charged amino acids at every third position. This region designated S4 is likely to be the voltage sensor of the protein. In the brain, HCN3 and HCN4 exhibit subcortical distribution mainly concentrated in the hypothalamus and thalamus, respectively.

Gene ID:

57657

### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200   |
|--------------------|--|
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |