



[Go to Product page](#)

Datasheet for ABIN5004533

anti-HES2 antibody (AA 1-100) (Alexa Fluor 680)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | HES2 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HES2 antibody is conjugated to Alexa Fluor 680 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human HES2 |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | HES2 |
| Alternative Name: | HES2 (HES2 Products) |

Target Details

| | |
|-------------|---|
| Background: | <p>Synonyms: bHLHb40, Class B basic helix loop helix protein 40, Hairy and enhancer of split 2 Drosophila, Transcription factor HES 2, HES2_HUMAN.</p> <p>Background: The Drosophila hairy and Enhancer of split genes encode basic helix-loop-helix (bHLH) transcriptional repressors that function in the Notch signaling pathway and control segmentation and neural development during embryogenesis. The mammalian homologues of Drosophila hairy and Enhancer of split are the HES gene family members, HES1-6, which also encode bHLH transcriptional repressors that regulate myogenesis and neurogenesis. The HES family members form a complex with TLE, the mammalian homologue of Groucho, and this interaction is mediated by the carboxy terminal WRPW motif of the HES proteins. The HES/TLE complex functions by directly binding to DNA, instead of interfering with activator proteins. Most HES family members, including HES1 and HES5, preferentially bind to the N box (CACNAG) as opposed to the E box (CANNTG). HES2 binds to both N and E box sites, while HES6 does not bind DNA. Rather, HES6 inhibits HES1 activity, thereby promoting transcription. HES1 and HES2 are expressed in a variety of adult and embryonic tissues. HES3 is expressed exclusively in cerebellar Purkinje cells, and HES5 is found solely in the nervous system. HES6 is produced in brain as well as in the limb buds of developing embryos.</p> |
|-------------|---|

| | |
|----------|-------|
| Gene ID: | 54626 |
|----------|-------|

Application Details

| | |
|--------------------|--|
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 |
|--------------------|--|

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|---------|--------|
| Format: | Liquid |
|---------|--------|

| | |
|----------------|---------|
| Concentration: | 1 µg/µL |
|----------------|---------|

| | |
|---------|---|
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
|---------|---|

| | |
|---------------|---------|
| Preservative: | ProClin |
|---------------|---------|

| | |
|--------------------|--|
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
|--------------------|--|

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months