antibodies .- online.com





anti-HIBADH antibody (AA 251-336) (Alexa Fluor 680)



\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	HIBADH	
Binding Specificity:	AA 251-336	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HIBADH antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HIBADH	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Cow,Sheep,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	HIBADH
Alternative Name:	HIBADH (HIBADH Products)

Target Details

9		
Background:	Synonyms: 3 hydroxy 2 methylpropanoate:NAD+ oxidoreductase, 3 hydroxyisobutyrate	
	dehydrogenase, 3 hydroxyisobutyrate dehydrogenase mitochondrial, EC 1.1.1.31, MGC40361,	
	NS5ATP1, 3HIDH_HUMAN.	
	Background: HIBADH is a 336 amino acid mitochondrial enzyme that catalyzes the NAD+-	
	$dependent, reversible\ oxidization\ of\ 3-Hydroxy is obutyrate\ to\ methylmal on at e\ semial dehyde,\ an$	
	intermediate of valine catabolism. The enzyme functions as a homodimer between a pH of 7.0	
	and 10.0, with optimal activity between 8.8 and 9.0. It was previously hypothesized that defects	
	in the gene encoding HIBADH may be the cause of 3-Hydroxyisobutyric aciduria, a rare disorder	
	that is characterized by a variety of clinical manifestations such as neurodevelopmental	
	problems and dysmorphic features. However, it was shown that HIBADH activity was equal in	
	patients with 3-Hydroxyisobutyric aciduria as compared with controls.	
Gene ID:	11112	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

12 months

Expiry Date: