

Datasheet for ABIN5004755

anti-HMGB1 antibody (AA 75-170) (AbBy Fluor® 750)



Go to Product page

| Overview | |
|----------|---|
| | ۰ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | HMGB1 | |
| Binding Specificity: | AA 75-170 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This HMGB1 antibody is conjugated to AbBy Fluor® 750 | |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human HMGB1 | | |
|-----------------------|---|--|--|
| Isotype: | IgG | | |
| Cross-Reactivity: | Human, Mouse, Rat | | |
| Predicted Reactivity: | Cow | | |
| Purification: | Purified by Protein A. | | |

Target Details

| Target: | HMGB1 |
|-------------------|------------------------|
| Alternative Name: | HMGB1 (HMGB1 Products) |

Target Details

| Background: | Synonyms: HMG1, HMG3, SBP-1, High mobility group protein B1, High mobility group protein 1, HMG-1, HMGB1 | | |
|---------------------|--|--|--|
| | Background: DNA binding proteins that associates with chromatin and has the ability to bend | | |
| | DNA. Binds preferentially single-stranded DNA. Involved in V(D)J recombination by acting as a | | |
| | cofactor of the RAG complex. Acts by stimulating cleavage and RAG protein binding at the 23 | | |
| | bp spacer of conserved recombination signal sequences (RSS). Heparin-binding protein that | | |
| | has a role in the extension of neurite-type cytoplasmic processes in developing cells (By | | |
| | similarity). | | |
| Gene ID: | 3146 | | |
| UniProt: | P09429 | | |
| Pathways: | p53 Signaling, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development, | | |
| | Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process, | | |
| | Toll-Like Receptors Cascades, Smooth Muscle Cell Migration, Inflammasome | | |
| Application Details | | | |
| Application Notes: | FCM 1:20-100 | | |
| | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| | | | |

| | 1 | l: |
|---|------|--------|
| - | ココロロ | urnen |
| | Hand | III IU |
| | | |

Expiry Date:

12 months