# antibodies -online.com





## anti-Indole 3 Acetic Acid antibody (Alexa Fluor 680)



Go to Product page

( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	Indole 3 Acetic Acid (IAA)
Reactivity:	Please inquire
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Indole 3 Acetic Acid antibody is conjugated to Alexa Fluor 680
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

#### **Product Details**

Immunogen:	KLH conjugated with IAA
Isotype:	IgG
Cross-Reactivity (Details):	Indole-3-Acetic Acid
Purification:	Purified by Protein A.

#### **Target Details**

Target:	Indole 3 Acetic Acid (IAA)	
Alternative Name:	IAA (IAA Products)	
Target Type:	Chemical	
Background:	Synonyms: Indole-3-Acetic Acid, indole-3-acetic acid_indol- yl-3-acetic acid.	

#### **Target Details**

Background: Indole-3-acetic acid, also known as IAA, is a heterocyclic compound that is an phytohormones called auxins. This colourless solid is probably the most important plant auxin. The molecule is derived from indole, containing a carboxymethyl group (acetic acid). IAA has many different effects, as all auxins do, such as inducing cell elongation and cell division with all subsequent results for plant growth and development. There are less expensive and metabolically stable synthetic auxin analogs on the market for use in horticulture, such as indole-3-butyric acid (IBA) and 1-naphthaleneacetic acid (NAA).

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	