antibodies -online.com





anti-Indole 3 Acetic Acid antibody (Alexa Fluor 750)



Go to Product page

/ N	1	-	K	/1	01	A /
0	1//	\vdash	1 \	/ I	-	\/\/
\sim	٧.	\sim	1 V		\sim	v v

Quantity:	100 μL	
Target:	Indole 3 Acetic Acid (IAA)	
Reactivity:	Please inquire	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Indole 3 Acetic Acid antibody is conjugated to Alexa Fluor 750	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated with IAA
Isotype:	IgG
Cross-Reactivity (Details):	Indole-3-Acetic Acid
Purification:	Purified by Protein A.

Target Details

Target:	Indole 3 Acetic Acid (IAA)	
Alternative Name:	IAA (IAA Products)	
Target Type:	Chemical	
Background:	Synonyms: Indole-3-Acetic Acid, indole-3-acetic acid_indol- yl-3-acetic acid.	

Target Details

Background: Indole-3-acetic acid, also known as IAA, is a heterocyclic compound that is an phytohormones called auxins. This colourless solid is probably the most important plant auxin. The molecule is derived from indole, containing a carboxymethyl group (acetic acid). IAA has many different effects, as all auxins do, such as inducing cell elongation and cell division with all subsequent results for plant growth and development. There are less expensive and metabolically stable synthetic auxin analogs on the market for use in horticulture, such as indole-3-butyric acid (IBA) and 1-naphthaleneacetic acid (NAA).

Application Details

Storage:

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration: 1 μg/μL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months

-20 °C