

Datasheet for ABIN5005047

anti-IDH3A antibody (AA 281-366) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IDH3A
Binding Specificity:	AA 281-366
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IDH3A antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IDH3A
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IDH3A
Alternative Name:	IDH3A (IDH3A Products)

Target Details

Background:	Synonyms: Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial, Isocitric dehydrogenase subunit alpha, NAD(+)-specific ICDH subunit alpha, IDH3A Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. IDH3A is the alpha subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase.
-------------	---

Gene ID:	3419
----------	------

UniProt:	P50213
----------	------------------------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	1 µg/µL
----------------	---------

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

Preservative:	ProClin
---------------	---------

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Handling

Expiry Date: 12 months