

Datasheet for ABIN5005127

**anti-IGF2 antibody (AA 25-51) (AbBy Fluor® 750)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	IGF2
Binding Specificity:	AA 25-51
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGF2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IGF II
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	IGF2
Alternative Name:	IGF 2 ( <a href="#">IGF2 Products</a> )

## Target Details

---

Background:	Synonyms: IGF-II, PP9974, C11orf43, Insulin-like growth factor II, Somatomedin-A, IGF2, PP1446 Background: The insulin-like growth factors possess growth-promoting activity. In vitro, they are potent mitogens for cultured cells. IGF-II is influenced by placental lactogen and may play a role in fetal development. Preptin undergoes glucose-mediated co-secretion with insulin, and acts as physiological amplifier of glucose-mediated insulin secretion. Exhibits osteogenic properties by increasing osteoblast mitogenic activity through phosphoactivation of MAPK1 and MAPK3.
Gene ID:	3481
UniProt:	<a href="#">P01344</a>
Pathways:	<a href="#">Hormone Activity</a> , <a href="#">Regulation of Hormone Metabolic Process</a> , <a href="#">Regulation of Hormone Biosynthetic Process</a> , <a href="#">Regulation of Carbohydrate Metabolic Process</a> , <a href="#">Activated T Cell Proliferation</a>

## Application Details

---

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months