antibodies .- online.com





anti-IL17C antibody (AA 101-197) (Alexa Fluor 750)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	IL17C	
Binding Specificity:	AA 101-197	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IL17C antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IL-17C
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IL17C
Alternative Name:	IL-17C (IL17C Products)

Target Details

•		
Background:	Synonyms: CX2, IL-17C, Interleukin-17C, Cytokine CX2, IL17C, UNQ561/PR01122	
	Background: Cytokine that plays a crucial role in innate immunity of the epithelium, including to	
	intestinal bacterial pathogens, in an autocrine manner. Stimulates the production of	
	antibacterial peptides and proinflammatory molecules for host defense by signaling through	
	the NF-kappa-B and MAPK pathways. Acts synergically with IL22 in inducing the expression of	
	antibacterial peptides, including S100A8, S100A9, REG3A and REG3G. Synergy is also observed	
	with TNF and IL1B in inducing DEFB2 from keratinocytes. Depending on the type of insult, may	
	have both protective and pathogenic properties, either by maintaining epithelial homeostasis	
	after an inflammatory challenge or by promoting inflammatory phenotype. Enhanced	
	IL17C/IL17RE signaling may also lead to greater susceptibility to autoimmune diseases.	
Gene ID:	27189	
UniProt:	Q9P0M4	
Pathways:	Cellular Response to Molecule of Bacterial Origin	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	