antibodies -online.com





anti-CD51/CD61 antibody (Alexa Fluor 750)



Go to Product page

\sim			
	$ \vee \cap$	r\/I	ΘM

Quantity:	100 μL	
Target:	CD51/CD61 (ITGAV/ITGB3)	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CD51/CD61 antibody is conjugated to Alexa Fluor 750	
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)),	
	Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Integrin Alpha V + Beta 3	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	CD51/CD61 (ITGAV/ITGB3)	
Alternative Name:	Integrin Alpha V + Beta 3 (ITGAV/ITGB3 Products)	
Background:	Synonyms: CD51, MSK8, VNRA, VTNR, Integrin alpha-V, Vitronectin receptor subunit alpha, ITGAV, CD61, GP3A, BDPLT2, GPIIIa, BDPLT16, Integrin beta-3, Platelet membrane glycoprotein	
	IIIa, ITGB3	

Background: ITAGV encodes integrin alpha chain V. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. The I-domain containing integrin alpha V undergoes post-translational cleavage to yield disulfide-linked heavy and light chains, that combine with multiple integrin beta chains to form different integrins. Among the known associating beta chains (beta chains 1,3,5,6, and 8, 'ITGB1', 'ITGB3', 'ITGB5', 'ITGB6', and 'ITGB8'), each can interact with extracellular matrix ligands, the alpha V beta 3 integrin, perhaps the most studied of these, is referred to as the Vitronectin receptor (VNR). In addition to adhesion, many integrins are known to facilitate signal transduction. The ITGB3 protein product is the integrin beta chain beta 3. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. A given chain may combine with multiple partners resulting in different integrins. Integrin beta 3 is found along with the alpha IIb chain in platelets. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling.

Gene ID:

3690, 3685

UniProt:

P05106, P06756

Application Details

Application Notes:

FCM 1:20-100

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months