antibodies -online.com





anti-IQCG antibody (Alexa Fluor 680)



| () | 1/0 | r\ / I | 014 | |
|-----|-----|--------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|-------------------|--|--|
| Target: | IQCG | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This IQCG antibody is conjugated to Alexa Fluor 680 | |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human IQCG | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | IQCG | |
| Alternative Name: | IQCG (IQCG Products) | |
| Background: | Synonyms: IQ domain containing protein G, IQ mot containing G, DKFZp434B227, FLJ11667, FLJ23571, FLJ37775, IQCG_HUMAN. | |

Background: IQCG is a 443 amino acid protein containing one IQ domain. Widely distributed in

nature, the IQ domain forms an amphiphilic seven-turn -helix capable of binding calmodulin in a

Target Details

Ca2+-independent manner. The level of intracellular calcium is tightly regulated in all eukaryotic cells. A modest increase in this level can result in a myriad of physiological responses, most of which are mediated by calmodulin (CaM), the universal calcium sensor. In acute T-lymphoid/myeloid leukemia, IQCG forms a complex with Nup98, an O-linked glycoprotein and a component of the nuclear pore complex. NUP98-IQCG complex bind co-activators and/or co-repressors, which suggest a role in transcriptional regulation.Nup98-IQCG complex inhibits 32Dcl3 cell apoptosis induced by Arabinofuranosylcytosine (Ara-C) and partially blocks granulocyte differentiation induced by G-CSF. IQCG exists as two isoforms due to alternatively splicing events.

Gene ID:

84223

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--|--|
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |