antibodies -online.com





anti-ITM2C antibody (AA 3-88) (Alexa Fluor 750)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	ITM2C
Binding Specificity:	AA 3-88
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ITM2C antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ITM2C
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

Target Details

Target:	ITM2C
Alternative Name:	ITM2C (ITM2C Products)
Background: Synonyms: BRI3, BRICD2C, BRICHOS domain containing 2C, Cerebral protein 14, CT-BRIS	

Target Details	
	E25C, Integral membrane protein 2C, Integral membrane protein 3, Itm2c, ITM2C_HUMAN, ITM3 antibodyNPD018, Transmembrane protein BRI3. Background: Negative regulator of beta amyloid peptide production. May inhibit the processing of APP by blocking its access to alpha-and beta-secretase. Binding to the beta-secretase-cleaved APP C-terminal fragment is negligible, suggesting that ITM2C is a poor gamma-secretase cleavage inhibitor. May play a role in TNF-induced cell death and neuronal differentiation.
Gene ID:	81618
UniProt:	Q9NQX7
Application Details	
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

12 months

Storage:

Expiry Date:

Storage Comment: