antibodies -online.com





anti-RAPTOR antibody (C-Term)





Go to Product page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Overview		
Quantity:	0.1 mg	
Target:	RAPTOR	
Binding Specificity:	C-Term	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Application:	Western Blotting (WB), Immunofluorescence (IF), Enzyme Immunoassay (EIA)	
Product Details		
Immunogen:	Raptor (CT) polyclonal antibody was raised against a 16 amino acid peptide from near the carboxy-terminus of mouse Raptor.	
	Carboxy-terminus of mouse Naptor.	
Isotype:	IgG	
Cross-Reactivity (Details):	Species reactivity (tested):Mouse	
Purification:	Peptide affinity chromatography	
Target Details		
Target:	RAPTOR	
Alternative Name:	RAPTOR (RAPTOR Products)	
Background:	The mammalian Target of Rapamycin (TOR, also known as mTOR) is an evolutionarily conserved serine/threonine kinase that regulates cell growth and cell cycle through its ability to	

integrate signals from nutrient levels and growth factors (reviewed in 1). Rapamycin inhibits

TOR activity resulting in reduced cell growth and reduced rates of cell cycle and cell proliferation (reviewed in 2). Raptor (regulatory associated protein of TOR) is a TOR-binding protein essential for TOR signaling in vivo. It acts as a TOR scaffold protein whose binding by TOR substrates is necessary for effective TOR-catalyzed phosphorylation (3). These substrates include the ribosomal protein S6 kinase (RP S6K) and the eukaryotic initiation factor 4E binding protein 4EBP1, proteins necessary for cell growth and proliferation and responsive to nutrient and mitogen levels (4). Raptor binds these proteins through a common 5 amino acid TOR-signaling (TOS) motif, mutation of this motif prevents the TOR-dependent phosphorylation of these proteins (5). Synonyms: KIAA1303, P150 target of rapamycin TOR-scaffold protein, RPTOR, Regulatory-associated protein of mTOR

UniProt:

Q8K4Q0

Pathways:

Warburg Effect

Application Details

Application Notes:

ELISA. Western blot: 2 µg/mL. Immunocytochemistry.

Other applications not tested.


Optimal dilutions are dependent on conditions and should be determined by the user.


Restrictions:

For Research Use only

Handling

Buffer:	PBS containing 0.02 % sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer.	





Immunofluorescence

Image 1. Immunocytochemistry of Raptor in L1210 cells with this product at 10 μ g/ml.

Western Blotting

Image 2. Western blot analysis of Raptor in L1210 cell lysate with this product at (A) 2 and (B) 4 μ g/ml.