# antibodies .- online.com





## anti-KCNMB1 antibody (AA 10-80) (Alexa Fluor 750)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	KCNMB1
Binding Specificity:	AA 10-80
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNMB1 antibody is conjugated to Alexa Fluor 750
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCNMB1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

#### **Target Details**

Target:	KCNMB1
Alternative Name:	KCNMB1 (KCNMB1 Products)

Target Details	
Background:	Synonyms: Calcium-activated potassium channel subunit beta-1, KCNMB1, BK channel subunit
	beta-1, Charybdotoxin receptor subunit beta-1, K(VCA)beta-1, Maxi K channel subunit beta-1,
	Slo-beta-1, Slo-beta
	Background: Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel.
	Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to
	KCNMA1 channel diversity. Increases the apparent Ca2+/voltage sensitivity of the KCNMA1
	channel. It also modifies KCNMA1 channel kinetics and alters its pharmacological properties. It
	slows down the activation and the deactivation kinetics of the channel. Acts as a negative
	regulator of smooth muscle contraction by enhancing the calcium sensitivity to KCNMA1. Its
	presence is also a requirement for internal binding of the KCNMA1 channel opener
	dehydrosoyasaponin I (DHS-1) triterpene glycoside and for external binding of the agonist
	hormone 17-beta-estradiol (E2). Increases the binding activity of charybdotoxin (CTX) toxin to
	KCNMA1 peptide blocker by increasing the CTX association rate and decreasing the
	dissociation rate.

Gene ID:

3779

### Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1 .	
ш	land	linc	٩
	ıanıu	11111	J

Expiry Date:

12 months