antibodies .- online.com





anti-FERMT2 antibody (AA 352-420) (Alexa Fluor 750)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	FERMT2	
Binding Specificity:	AA 352-420	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FERMT2 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Kindlin 2/MIG2	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	FERMT2	
Alternative Name:	Kindlin 2/MIG2 (FERMT2 Products)	

Target Details

Background:	Synonyms: FLJ34213, FLJ44462, UNC112, FERM2_HUMAN, Fermitin family homolog 2,	
	Fermt2, KIND2, Kindlin-2, MIG-2, MIG2, Mitogen-inducible gene 2 protein, PH domain-containing	
	family C member 1, Pleckstrin homology domain containing family C with FERM domain	
	member 1, Pleckstrin homology domain-containing family C member 1, PLEKHC1, UNC112B.	
	Background: Mig-2 recruits migfilin to cell-matrix adhesions, while the interaction with filamin	
	mediates the association of migfilin with actin filaments. Together, Mig-2, migfilin and filamin	
	define a connection between cell matrix adhesions and the actin cytoskeleton and participate in	
	the orchestration of actin assembly and cell shape modulation. Mig-2 expression is	
	transcriptionally elevated in leiomyomas and could be involved in its hormone-mediated growth	
	of leiomyomas of the uterus. Expression of Mig-2 is ubiquitous, and it is found in numerous	
	tumor tissues.	
Gene ID:	10979	
Pathways:	Cell-Cell Junction Organization	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	