

Datasheet for ABIN5006130

anti-LEFTY1 antibody (AA 51-150) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	LEFTY1
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LEFTY1 antibody is conjugated to AbBy Fluor® 750
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Lefty
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	LEFTY1
Alternative Name:	Lefty (LEFTY1 Products)
Background:	Synonyms: Protein lefty-1, AI450052, Ebaf, Left right determination factor 1, left right

Target Details

determination factor, Left right determination factor B, left-right determination factor, LEFT-RIGHT DETERMINATION FACTOR 1, LEFTY1, left-right determination, factor B, LEFT-RIGHT DETERMINATION, FACTOR B, LEFTY B, LEFTB, LEFTY 1, Lefty, lefty-1, LEFTY1, LEFTY1, MOUSE, HOMOLOG OF antibody, LEFTYB, Protein lefty-B, RGD1561867, Stimulated by retinoic acid gene 3 protein, STRA3, TGF beta 4, TGF-beta-4, Tgfb4, Transforming growth factor beta 4, Transforming growth factor beta-4.

Background: During vertebrate embryogenesis, a left right axis is established. Secreted growth factors of the TGF beta family, including gene products derived from nodal, lefty 1 and lefty 2, play crucial roles in establishing left right asymmetries. TGF beta (Transforming growth factor beta) is a pleiotropic cytokine that regulates growth and differentiation of diverse types of cells. TGF beta actions are directed by ligand induced activation of TGF beta receptors. Complexes formed move into the nucleus, where they act as components of a transcriptional complex. Lefty, a novel member of the TGF beta superfamily, inhibits TGF beta signaling. Lefty acts to inhibit phosphorylation of Smad2 following activation of the TGF beta receptor. Lefty also inhibits events downstream from R Smad phosphorylation. Lefty provides a repressed state of TGF beta responsive genes. The Lefty family is comprised of Lefty 1 and Lefty 2 in mouse, and Lefty A and Lefty B in humans. Members of the TGF beta superfamily require processing for their activation. Cleavage is therefore an essential step for Lefty activation. Lefty is synthesized as a large inactive precursor (41 Kda) that must be endoproteolytically processed to release the bioactive polypeptide (28 kDa and 34 kDa forms). The 28 kDa form induces MAPK activity.

Gene ID: 10637

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months