# antibodies -online.com





# anti-LMCD1 antibody (AA 51-160) (Alexa Fluor 680)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | LMCD1  |
| Binding Specificity: | AA 51-160  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This LMCD1 antibody is conjugated to Alexa Fluor 680   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human LMCD1 |
|-----------------------|---|
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig,Horse,Rabbit                      |
| Purification:         | Purified by Protein A.                                    |

#### **Target Details**

| ranget betand     |  |  |
|-------------------|--|--|
| Target:           | LMCD1  |  |
| Alternative Name: | LMCD1 (LMCD1 Products)   |  |
| Background:       | Synonyms: Dyxin, LIM and cysteine rich domains 1, LIM and cysteine rich domains protein 1, |  |

LIM and cysteine-rich domains protein 1, LMCD1\_HUMAN.

Background: Transcriptional cofactor that restricts GATA6 function by inhibiting DNA-binding, resulting in repression of GATA6 transcriptional activation of downstream target genes.

Represses GATA6-mediated trans activation of lung- and cardiac tissue-specific promoters.

Inhibits DNA-binding by GATA4 and GATA1 to the cTNC promoter (By similarity). Plays a critical role in the development of cardiac hypertrophy via activation of calcineurin/nuclear factor of activated T cells signaling pathway.

## **Application Details**

| Restrictions:      | For Research Use only |
|--------------------|-----------------------|
|                    | IF(ICC) 1:50-200      |
|                    | IF(IHC-F) 1:50-200    |
| Application Notes: | IF(IHC-P) 1:50-200    |

### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |