antibodies .- online.com





anti-PPAP2C antibody (AA 151-250) (Alexa Fluor 750)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	PPAP2C	
Binding Specificity:	AA 151-250	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PPAP2C antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human VANGL1/LPP2	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	PPAP2C	
Alternative Name:	LPP2 (PPAP2C Products)	
Background: Synonyms: Loop tail protein 2 homolog, Loop-tail protein 2 homolog, LPP2, MGC5338, STB2		

STBM2, Strabismus 2, Van Gogh like protein 1, Van Gogh-like protein 1, vang like 1 van gogh, Drosophila, Vang like protein 1, Vang-like protein 1, VANG1_HUMAN, VANGL 1, Vangl1.

Background: The Vang family of proteins are integral membrane proteins that are homologues of the Drosophila tissue polarity gene strabismus. The gene encoding for Van Gogh-like protein 1 (Vangl1), also designated Strabismus 2 (STB2), localizes to human chromosome 1p11-p13.1.

Van Gogh-like protein 2 (Vangl2), also designated Strabismus 1 (STB1), localizes to chromosome 1q22-q23. Vangl1 is expressed primarily in testis and ovary, but is also expressed in gastric and pancreatic cancer. Vangl proteins play a key developmental role in establishing planar cell polarity (PCP) and in regulating convergent extension (CE) movements during embryogenesis. Vangl1 and Vangl2 are both downregulated in several cancer cell lines and primary tumors.

Gene ID:

8612

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	