antibodies -online.com





anti-LRFN2 antibody (AA 401-500) (Alexa Fluor 750)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	LRFN2	
Binding Specificity:	AA 401-500	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LRFN2 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LRFN2/SALM1	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	LRFN2
Alternative Name:	LRFN2/SALM1 (LRFN2 Products)

Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		\sim	<i>.</i>	ч.

Synonyms: leucine rich repeat and fibronectin type III domain containing 2, RP11-535K1.2, fibronectin type III, immunoglobulin and leucine rich repeat domains 2 antibody FIGLER2, Leucine-rich repeat and fibronectin type-III domain-containing protein 2, LRFN2, LRFN2_HUMAN, SALM1, Synaptic adhesion-like molecule 1.

Background: LRFN2 is a 789 amino acid single-pass type I membrane protein belonging to the LRFN family. Encoded by a gene that maps to human chromosome 6p21.2, LRFN2 is moderately expressed in brain, spleen and testis. LRFN2 contains one fibronectin type-III domain, one Ig-like (immunoglobulin-like) domain and six LRR (leucine-rich) repeats. LRFN2 promotes neurite outgrowth in hippocampal neurons, enhances cell surface expression of two NMDA receptor subunits, NMDA1 and NMDA?1, and may play a role in redistributing PSD-95 to cell periphery. LRFN2 forms heteromeric complexes with LRFN1, LRFN3, LRFN4 and LRFN5, and is capable of forming homomeric complexes, but not across cell junctions.

Pathways:

Synaptic Membrane

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months