

Datasheet for ABIN5006304

anti-LRP5 antibody (AA 501-600) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	LRP5
Binding Specificity:	AA 501-600
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRP5 antibody is conjugated to AbBy Fluor® 750
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LRP5
Isotype:	IgG
Specificity:	This antibody will recognize LRP6 due to high sequence similarity.
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	LRP5
---------	------

Target Details

Alternative Name:	LRP5 (LRP5 Products)
Background:	Synonyms: HBM, LR3, OPS, EVR1, EVR4, LRP7, OPPG, BMND1, LRP-5, OPTA1, VBCH2, Low-density lipoprotein receptor-related protein 5, LRP5 Background: Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalsomes. Cell-surface coreceptor of Wnt/beta-catenin signaling, which plays a pivotal role in bone formation. The Wnt-induced Fzd/LRP6 coreceptor complex recruits DVL1 polymers to the plasma membrane which, in turn, recruits the AXIN1/GSK3B-complex to the cell surface promoting the formation of signalsomes and inhibiting AXIN1/GSK3-mediated phosphorylation and destruction of beta-catenin. Appears be required for postnatal control of vascular regression in the eye. Required for posterior patterning of the epiblast during gastrulation.
Gene ID:	4041
UniProt:	O75197
Pathways:	WNT Signaling , Stem Cell Maintenance , Positive Regulation of fat Cell Differentiation

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months