

Datasheet for ABIN5006385

anti-LYN antibody (AA 201-300) (AbBy Fluor® 680)



Go to Product page

_				
()	1//	rv	IO	Λ/
()	VC	. I V	1	v v

Quantity:	100 μL	
Target:	LYN	
Binding Specificity:	AA 201-300	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LYN antibody is conjugated to AbBy Fluor® 680	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Lyn	
Isotype:	IgG	
Specificity:	This protein is known to be similar in amino acid sequence to HCK. Therefore, cross-reactivity with this homologous protein may be observed	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Background: Synonyms: JTK8, p53Lyn, p56Lyn, Tyrosine-protein kinase Lyn, Lck/Yes-related novel protein tyrosine kinase, V-yes-1 Yamaguchi sarcoma viral related oncogene homolog, LYN Background: This gene encodes a tyrosine protein kinase, which maybe involved in the regulation of mast cell degranulation, and erythroid differentiation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq. Jul 2011] Gene ID: 4067 UniProt. P07948 Pathways: Fo epsilon Receptor Signaling Pathway, Hormone Transport, Response to Growth Hormone Simulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Target:	LYN	
tyrosine kinase, V-yes-1 Yarnaguchi sarcoma viral related oncogene homolog, LYN Background: This gene encodes a tyrosine protein kinase, which maybe involved in the regulation of mast cell degranulation, and erythroid differentiation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq. Jul 2011] Gene ID: 4067 UniProt: P07948 Pathways: Fo-epsilon Receptor Signaling Pathway, Hormone Transport, Response to Growth Hormone Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FOM 1:20-100 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Alternative Name:	Lyn (LYN Products)	
transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2011] Gene ID: 4067 UniProt: P07948 Pathways: Fc-epsilon Receptor Signaling Pathway, Hormone Transport, Response to Growth Hormone Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Background:	tyrosine kinase, V-yes-1 Yamaguchi sarcoma viral related oncogene homolog, LYN Background: This gene encodes a tyrosine protein kinase, which maybe involved in the	
UniProt: Pathways: Fo-epsilon Receptor Signaling Pathway, Hormone Transport, Response to Growth Hormone Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 am 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		transcript variants encoding different isoforms have been found for this gene. [provided by	
Pathways: Fc-epsilon Receptor Signaling Pathway, Hormone Transport, Response to Growth Hormone Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Gene ID:	4067	
Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling, Integrin Complex, BCR Signaling Application Details Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	UniProt:	P07948	
Application Notes: FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Pathways:	Stimulus, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events,	
IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Application Details		
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200	
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Restrictions:	For Research Use only	
Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Handling		
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Format:	Liquid	
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Concentration:	1 μg/μL	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
handled by trained staff only.	Preservative:	ProClin	
Storage: -20 °C	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
	Storage:	-20 °C	

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months