# antibodies .- online.com





## anti-MAP9 antibody (AbBy Fluor® 750)



( )	1/0	r\ / I	014	
( )	ve	I V I	-v	V

Overview	
Quantity:	100 μL
Target:	MAP9
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP9 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human MAP-9
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	MAP9
Alternative Name:	MAP-9 (MAP9 Products)
Background:	Synonyms: ASAP, Aster associated protein, MAP9, MAP 9, microtubule associated protein 9, MAP9_HUMAN.
	Background: Microtubules, the primary component of the cytoskeletal network, interact with
	proteins called microtubule-associated proteins (MAPs).MAP9 is a microtubule-associated

protein required for spindle function, mitotic progression, and cytokinesis. The microtubule-associated proteins can be divided into two groups, structural and dynamic. The MAP proteins function to stimulate tubulin assembly, enhance microtubule stability, influence the spatial distribution of microtubules within cells and utilize microtubule polarity to translocate cellular components. MAP-9 (microtubule-associated protein 9), also known as ASAP, is a 647 amino acid cytoplasmic protein that is constitutively expressed during the cell cycle. MAP-9 localizes to microtubules in interphase, associates with the mitotic spindle during mitosis and localizes to the central body during cytokinesis. Involved in organization of the bipolar mitotic spindle, MAP-9 is required for bipolar spindle assembly, mitosis progression and cytokinesis. MAP-9 may be involved in stabilizing interphase microtubules. Two isoforms of MAP-9 are produced due to alternative splicing events.

Gene ID:

79884

### Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

 Concentration:
 1 μg/μL

 Buffer:
 Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

 Preservative:
 ProClin

 Precaution of Use:
 This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

 Storage:
 -20 °C

 Storage Comment:
 Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

 Expiry Date:
 12 months