

## Datasheet for ABIN5006652

## anti-M-CSF/CSF1 antibody (AA 201-300) (AbBy Fluor® 750)



Go to Product page

	er		

Quantity:	100 μL		
Target:	M-CSF/CSF1 (CSF1)		
Binding Specificity:	AA 201-300		
Reactivity:	Human, Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This M-CSF/CSF1 antibody is conjugated to AbBy Fluor® 750		
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human MCSF		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse		
Predicted Reactivity:	Rat,Dog,Cow,Pig,Rabbit		
Purification:	Purified by Protein A.		
Target Details			
Target:	M-CSF/CSF1 (CSF1)		
Alternative Name:	MCSF (CSF1 Products)		

## **Target Details**

Synonyms: MCSF, CSF-1, Macrophage colony-stimulating factor 1, M-CSF, Lanimostim, CSF1		
Background: Cytokine that plays an essential role in the regulation of survival, proliferation and		
differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as		
macrophages and monocytes. Promotes the release of proinflammatory chemokines, and		
thereby plays an important role in innate immunity and in inflammatory processes. Plays an		
important role in the regulation of osteoclast proliferation and differentiation, the regulation of		
bone resorption, and is required for normal bone development. Required for normal male and		
female fertility. Promotes reorganization of the actin cytoskeleton, regulates formation of		
membrane ruffles, cell adhesion and cell migration. Plays a role in lipoprotein clearance.		
1435		
P09603		
RTK Signaling		
IF(IHC-P) 1:50-200		
IF(IHC-F) 1:50-200		
IF(ICC) 1:50-200		
For Research Use only		
Liquid		
1 μg/μL		
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
50 % Glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
handled by trained staff only.		
-20 °C		
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
12 months		
• • • • • • • • • • • • • • • • • • •		