# antibodies -online.com





# anti-LY86 antibody (AA 65-162) (AbBy Fluor® 750)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | LY86   |
| Binding Specificity: | AA 65-162  |
| Reactivity:          | Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This LY86 antibody is conjugated to AbBy Fluor® 750  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human LY-86 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Rat   |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Pig,Rabbit                            |
| Purification:         | Purified by Protein A.                                    |

## **Target Details**

| Target:           | LY86                |
|-------------------|---------------------|
| Alternative Name: | MD1 (LY86 Products) |

## **Target Details**

| Background:         | Synonyms: BPAD, dJ80N2.1, LY86, LY-86, LY 86, Lymphocyte antigen 86, Lymphocyte antigen          |  |
|---------------------|--|--|
|                     | 86 precursor, Major affective disorder 1, MD-1, MD1, MD 1, MD1 protein, MD1 RP105                |  |
|                     | associated, MD1 RP105 associated, MMD-1, RP1-80N2.1.   |  |
|                     | Background: MD1 is a 28 kDa molecule that is associated at the cell surface with RP105           |  |
|                     | (CD180). RP105 is a leucine-rich repeat (LRR) molecule that is expressed on B lymphocytes. It    |  |
|                     | was first identified by a mAb that protects spleen B cells from irradiation-induced apoptosis.   |  |
|                     | LRR proteins, such as Toll receptors, have a role in innate immunity. MD1 is primarily expressed |  |
|                     | by B cells, dendritic cells and monocytes, and promotes the recognition of, and subsequent       |  |
|                     | signaling by LPS in the innate immune system. MD1 does not have any homology to other            |  |
|                     | molecules. MD1 when expressed alone behaves as a secretory protein. Expression of MD1 in a       |  |
|                     | cell increases the expression of RP105.  |  |
| Gene ID:            | 9450   |  |
| Pathways:           | Toll-Like Receptors Cascades   |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and      |  |
|                     | 50 % Glycerol.   |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be              |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |