



[Go to Product page](#)

Datasheet for ABIN5006683  
**anti-MEF2B antibody (Alexa Fluor 680)**

### Overview

|              |   |
|--------------|---|
| Quantity:    | 100 µL  |
| Target:      | MEF2B   |
| Reactivity:  | Human, Mouse, Rat   |
| Host:        | Rabbit  |
| Clonality:   | Polyclonal  |
| Conjugate:   | This MEF2B antibody is conjugated to Alexa Fluor 680                            |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                   |   |
|-------------------|---|
| Immunogen:        | KLH conjugated synthetic peptide derived from human MEF2B |
| Isotype:          | IgG   |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Purified by Protein A.                                    |

### Target Details

|                   |  |
|-------------------|--|
| Target:           | MEF2B  |
| Alternative Name: | MEF2B ( <a href="#">MEF2B Products</a> )   |
| Background:       | Synonyms: PRDM3, MADS box transcription enhancer factor 2, MDS1 EVI1, Mef2b, MEF2B_HUMAN, MEF-2B, myocyte enhancer factor 2B, myocyte specic enhancer factor 2B, Myocyte-specific enhancer factor 2B, RSRFR2, serum response factor like protein 2, Serum response factor-like protein 2, XMEF2. |

## Target Details

---

Background: The myocyte enhancer factor-2 (MEF-2) family of transcription factors associate with co-repressors or co-activators to regulate development and function of T cells, neuronal cells, and muscle cells. Four family members, termed MEF-2A, -2B, -2C, and -2D, arise from alternatively spliced transcripts. These members bind as homo- and heterodimers to the MEF-2 site in the promoter region of affected genes. Differential regulation in the expression of the four transcripts implies functional distinction for each during embryogenesis and development. The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including MyoD, myogenin, Myf-5, and MRF4, are one class of identified factors. The MEF-2 family represents a second class of DNA binding regulatory proteins. Each of these proteins binds to the MEF-2 target DNA sequence present in the regulatory regions of many muscle-specific genes.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months