# antibodies .- online.com





# anti-MFSD2A antibody (AA 331-430) (Alexa Fluor 750)



| $C_0 + c$ | Produc | +      |
|-----------|--------|--------|
|           | Promi  | บ กลกย |
|           |        |        |

| ( ) | 1/0 | r\ / I | 014 |   |
|-----|-----|--------|-----|---|
| ( ) | ve  | I V I  | -v  | V |
|     |     |        |     |   |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | MFSD2A  |
| Binding Specificity: | AA 331-430  |
| Reactivity:          | Human, Mouse, Rat   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This MFSD2A antibody is conjugated to Alexa Fluor 750   |
| Application:         | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human MFSD2A |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Mouse, Rat  |
| Predicted Reactivity: | Dog,Cow,Pig,Horse,Chicken,Rabbit                           |
| Purification:         | Purified by Protein A.                                     |

## **Target Details**

| Target:           | MFSD2A                   |
|-------------------|--------------------------|
| Alternative Name: | MFSD2A (MFSD2A Products) |

#### Target Details

| Bac | kar | 'nΙ    | ın       | Ч. |
|-----|-----|--------|----------|----|
| Duo |     | $\sim$ | <i>.</i> | ч. |

Synonyms: NLS1, MFSD2, Sodium-dependent lysophosphatidylcholine symporter 1, Sodium-dependent LPC symporter 1, Major facilitator superfamily domain-containing protein 2A, MFSD2A, HMFN0656, PP9177, UNQ300/PRO341

Background: Sodium-dependent lysophosphatidylcholine (LPC) symporter, which plays an essential role for blood-brain barrier formation and function. Specifically expressed in endothelium of the blood-brain barrier of micro-vessels and transports LPC into the brain. Transport of LPC is essential because it constitutes the major mechanism by which docosahexaenoic acid (DHA), an omega-3 fatty acid that is essential for normal brain growth and cognitive function, enters the brain. Transports LPC carrying long-chain fatty acids such LPC oleate and LPC palmitate with a minimum acyl chain length of 14 carbons. Does not transport docosahexaenoic acid in unesterified fatty acid. Specifically required for blood-brain barrier formation and function, probably by mediating lipid transport. Not required for central nervous system vascular morphogenesis (By similarity). Acts as a transporter for tunicamycin, an inhibitor of asparagine-linked glycosylation. In placenta, acts as a receptor for ERVFRD-1/syncytin-2 and is required for trophoblast fusion (PubMed:18988732).

Gene ID: 84879

UniProt: Q8NA29

#### **Application Details**

Application Notes:

FCM 1:20-100

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

| Format:            | Liquid  |
|--------------------|---|
| Concentration:     | 1 μg/μL   |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                         |

## Handling

|                  | handled by trained staff only.  |
|------------------|---|
| Storage:         | -20 °C  |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date:     | 12 months   |