

Datasheet for ABIN5006840

anti-MIB1 antibody (AA 101-200) (AbBy Fluor® 680)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MIB1
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MIB1 antibody is conjugated to AbBy Fluor® 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Mib1/Mindbomb
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	MIB1
Alternative Name:	Mib1/Mindbomb (MIB1 Products)
Background:	Synonyms: DAPK-interacting protein 1, Dip 1, DIP-1, Dip1, E3 ubiquitin protein ligase MIB 1, E3

Target Details

ubiquitin protein ligase MIB1, E3 ubiquitin-protein ligase mib1, mib1, MIB1_HUMAN, Mind bomb homolog 1, Zinc finger ZZ type with ankyrin repeat domain protein 2.

Background: MIB1 is a 1006 amino acid E3 ubiquitin ligase that activates the Notch ligand, Delta. MIB1 ubiquitinates Delta by binding to its intracellular domain, leading to the endocytosis and eventual degradation of the Delta receptor, which, paradoxically, results in the up-regulation of receptor activity and enhances Notch signaling. MIB1 also interacts with DAPK, a protein that plays an important role in the regulation of apoptosis. Ubiquitination of DAPK leads to inhibition of caspase-dependent apoptosis, therefore it is likely that overexpression of MIB1 can lead to tumor growth. Although it seems to be widely expressed at low levels, MIB1 is expressed at highest concentrations in the CNS and ovary. Both DAPK and MIB1 are overexpressed in epileptic brain tissue, suggesting that they probably cooperate as regulators of neuronal death in epilepsy.

Gene ID: 57534

Pathways: [SARS-CoV-2 Protein Interactome](#), [The Global Phosphorylation Landscape of SARS-CoV-2 Infection](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months