
antibodies - online.com







anti-MCPH1 antibody (AA 11-110) (Alexa Fluor 750)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	MCPH1
Binding Specificity:	AA 11-110
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MCPH1 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Microcephalin 1/BRIT1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Dog
Purification:	Purified by Protein A.

Target Details

- Target Details	
Target:	MCPH1
Alternative Name:	Microcephalin 1/BRIT1 (MCPH1 Products)

Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		\sim	<i>.</i>	ч.

Synonyms: BRCT repeat inhibitor of TERT expression 1, BRIT 1, FLJ12847, Hypothetical protein FLJ12847, MCPH 1, MCPH1_HUMAN, MCT antibody Microcephalin 1, Microcephalin-1, Microcephaly primary autosomal recessive 1.

Background: Microcephalin modulates brain size and has been proliferating under strong positive selection for several thousand years, although the nature of the positive selection is poorly understood. Human Microcephalin contains three BRCA1 C-terminal (BRCT) domains and shares 57 % identity with its mouse ortholog, the most conserved regions being BRCT domains where there is 80 % identity. Predominant expression of human Microcephalin is observed in fetal brain, liver and kidney tissues and is expressed during neurogenesis in mice. Microcephalin displays significantly higher rates of protein evolution in primates than in rodents, this trend is most noticeable for the subset of genes associated with nervous system development. Microcephalin has a very young, single nucleotide, polymorphism haplotype associated with modern humans, this gene is presumably still evolving in Homo sapiens. It functions in DNA damage response and regulation of cell cycle checkpoints.

Pathways:

Stem Cell Maintenance

Application Details

Λnn	lication	Motac.
\neg	псанон	TAUTES.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months