# antibodies -online.com





## anti-MNAT1 antibody (AA 51-150) (Alexa Fluor 680)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	MNAT1	
Binding Specificity:	AA 51-150	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MNAT1 antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MNAT1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target:	MNAT1
Alternative Name:	MNAT1 (MNAT1 Products)

### Target Details

Background:	Synonyms: MAT1, TFB3, CAP35, RNF66, CDK-activating kinase assembly factor MAT1,	
	CDK7/cyclin-H assembly factor, Cyclin-G1-interacting protein, Menage a trois, RING finger	
	protein 66, RING finger protein MAT1, p35, p36, MNAT1	
	Background: Stabilizes the cyclin H-CDK7 complex to form a functional CDK-activating kinase	
	(CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and	
	CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIH basal transcription	
	factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal	
	domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and	
	elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA	
	polymerase II.	
Gene ID:	4331	
UniProt:	P51948	
Pathways:	Cell Division Cycle, Mitotic G1-G1/S Phases, M Phase	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	