

Datasheet for ABIN5007120

anti-MTOR antibody (AA 2436-2492) (AbBy Fluor® 680)



Go to Product page

()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Quantity:	100 μL
Target:	MTOR (mTOR)
Binding Specificity:	AA 2436-2492
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTOR antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human mTOR	
Isotype:	IgG	
Cross-Reactivity:	Chicken, Cow, Fish, Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Sheep,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	MTOR (mTOR)
Alternative Name:	mTOR/ FRAP (mTOR Products)

Target Details

Bac	kgr	ou	nd	:

Synonyms: Serine/threonine-protein kinase mTOR, FK506-binding protein 12-rapamycin complex-associated protein 1, FKBP12-rapamycin complex-associated protein, Mammalian target of rapamycin, mTOR, Mechanistic target of rapamycin, Rapamycin and FKBP12 target 1, Rapamycin target protein 1, MTOR, FRAP, FRAP1, FRAP2, RAFT1, RAPT1

Background: The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This protein acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. The ANGPTL7 gene is located in an intron of this gene. [provided by RefSeq, Sep 2008]

Gene ID:

2475

UniProt:

P42345

Pathways:

PI3K-Akt Signaling, RTK Signaling, AMPK Signaling, Interferon-gamma Pathway, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Regulation of Actin Filament Polymerization, Regulation of Muscle Cell Differentiation, Regulation of Cell Size, Skeletal Muscle Fiber Development, Regulation of Carbohydrate Metabolic Process, Autophagy, CXCR4-mediated Signaling Events, BCR Signaling, Warburg Effect

Application Details

Δn	plicati	on N	Intac
ΑD	piicati	on n	10162

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months