

Datasheet for ABIN5007257
anti-NAB2 antibody (AbBy Fluor® 750)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NAB2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NAB2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NAB2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NAB2
Alternative Name:	NAB2/ MADER (NAB2 Products)
Background:	Synonyms: EGR 1 binding protein 2, EGR-1-binding protein 2, EGR1 binding protein 2, MADER, Melanoma associated delayed early response protein, Melanoma-associated delayed early

Target Details

response protein, MGC75085, Nab 2, nab2, NAB2_HUMAN, NGFI A binding protein 2 EGR1 binding protein 2, NGFI A binding protein 2, NGFI-A-binding protein 2, Protein MADER.

Background: Transcriptional control is in part regulated by interactions between DNA-bound transcription factors, such as Egr-1/NGFI-A, and coregulatory proteins, such as NAB (for NGFI-A-binding proteins). The evolutionarily conserved NAB proteins, NAB1 and NAB2 are corepressors of EGF-1/NGFI-A. Both NAB1 and NAB2 contain an amino terminal NAB conserved domain 1 (NCB1), which is required for binding NGFI-A, and a carboxy terminal NCD2 domain, which is responsible for the repressor function of NAB proteins. NAB2 is principally localized in the nucleus and may play a role in the downregulation of NGFI-A activity as well as in controlling fundamental processes such as cell division, differentiation and apoptosis. NAB2 localizes to chromosome 12q13.3-14.1, a region that is rearranged in several solid tumors, lipomas, and liposarcomas.

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months