



[Go to Product page](#)

Datasheet for ABIN5007444

anti-NETO2 antibody (AA 151-250) (Alexa Fluor 750)

Overview

Quantity:	100 µL
Target:	NETO2
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NETO2 antibody is conjugated to Alexa Fluor 750
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NETO2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig
Purification:	Purified by Protein A.

Target Details

Target:	NETO2
Alternative Name:	NETO2 (NETO2 Products)
Background:	Synonyms: Brain specic transmembrane protein, Brain specic transmembrane protein

Target Details

containing 2 CUB and 1 LDL receptor class A domains protein 2, Brain-specific transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2, BTCL2, FLJ10430, FLJ14724, FLJ90456, NEOT2, NETO 2, Neto2, NETO2_HUMAN, Neuropilin NRP and tolloid TLL like 2, Neuropilin and tolloid like protein 2, Neuropilin and tolloid-like protein 2.

Background: NETO2 is a 525 amino acid single-pass type I membrane protein that contains two CUB domains and one LDL-receptor class A domain. Expressed as multiple alternatively spliced isoforms, NETO2 is thought to play a role in the development and maintenance of neuronal circuitry, possibly playing a role in proper brain function. The gene encoding NETO2 maps to human chromosome 16, which is associated with a variety of genetic disorders, encodes over 900 genes and comprises nearly 3 % of the human genome. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is associated with chromosome 16, as is Crohn's disease, which is a gastrointestinal inflammatory condition.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months