antibodies -online.com





anti-Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1) (AA 34-111) antibody (Alexa Fluor 680)



Go to Product page

| \sim | | | | | |
|--------|-----|----|-----|-------------|----|
| | 1/6 | ⊃r | \/I | \triangle | ٨/ |

| - Overview | | |
|----------------------|---|--|
| Quantity: | 100 μL | |
| Target: | Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1) | |
| Binding Specificity: | AA 34-111 | |
| Reactivity: | Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | Alexa Fluor 680 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |
| Product Details | | |

| Immunogen: | KLH conjugated synthetic peptide derived from human NADE/NGFRAP1 | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Rat | |
| Predicted Reactivity: | Human,Mouse,Dog,Cow,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1) |
|-------------------|--|
| Alternative Name: | NGFRAP1 (NGFRAP1 Products) |

Target Details

| Background: | Synonyms: BEX3, BEX3_HUMAN, Brain expressed X linked 3 mouse homolog, Brain-expressed | |
|---------------------|--|--|
| | X-linked protein 3, Nerve growth factor receptor TNFRSF16 associated protein 1, Nerve growth | |
| | factor receptor associated protein 1, Nerve growth factor receptor-associated protein 1, | |
| | NGFRAP1, Ovarian granulosa cell 13.0 kDa protein HGR74, Ovarian granulosa cell protein, | |
| | p75NTR associated cell death executor, p75NTR-associated cell death executor, Protein BEX3. | |
| | Background: May be a signaling adapter molecule involved in p75NTR-mediated apoptosis | |
| | induced by NGF. Plays a role in zinc-triggered neuronal death (By similarity). May play an | |
| | important role in the pathogenesis of neurogenetic diseases. Tissue specificity: Found in | |
| | ovarian granulosa cells, testis, prostate and seminal vesicle tissue. High levels also detected in | |
| | liver. | |
| Gene ID: | 27018 | |
| Pathways: | Neurotrophin Signaling Pathway, Positive Regulation of Endopeptidase Activity | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |