

Datasheet for ABIN5007724

anti-NME4 antibody (AA 21-120) (AbBy Fluor® 680)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NME4
Binding Specificity:	AA 21-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NME4 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NME4/nm23-H4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NME4
Alternative Name:	NME4/nm23-H4 (NME4 Products)

Target Details

Background:	Synonyms: mitochondrial, NDK, NDKM_HUMAN, NDP kinase, NDP kinase, mitochondrial, NDPKD, nm23 H4, nm23-H4, NM23D, Nm23M4, NME4, Non metastatic cells 4 protein expressed in, Nucleoside diphosphate kinase D, Nucleoside diphosphate kinase, mitochondrial. Background: The nm23 gene (Metastasis inhibition factor nm23), a potential suppressor of metastasis, is expressed at a much lower level in highly metastatic cells than in cells with lower metastatic potential. Based on sequence analysis, nm23, also designated nucleoside diphosphate kinase A (NDK A) or Tumor metastatic process-associated protein, appears to be highly related to nucleotide diphosphate kinases (NDP). NDP kinases A and B are identical to two isotypes of human nm23 homologs, nm23-H1 and nm23-H2, respectively. nm23-H2 is also identical in sequence to PuF, a transcription factor that binds to nuclease-hypersensitive elements at positions 142 to 115 of the human c-Myc promoter. nm23-H3 and nm23-H4 are important for the synthesis of nucleoside triphosphates and may play a role in apoptosis induction and hematopoiesis. nm23-H4 localizes to the mitochondrial intermembrane space and is widely expressed, with higher levels detected in prostate, heart, liver, small intestine, and skeletal muscle tissues. Low amounts of nm23-H4 are observed in the brain and in blood leukocytes.
Pathways:	Nucleotide Phosphorylation , Ribonucleoside Biosynthetic Process

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months