antibodies.com

## Datasheet for ABIN5007746 anti-NOB1 antibody (Alexa Fluor 680)



Overview

Overview	
Quantity:	100 μL
Target:	NOB1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOB1 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human NOB1
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	NOB1
Alternative Name:	NOB1 (NOB1 Products)
Background:	Synonyms: Adenocarcinoma antigen recognized by T lymphocytes 4, ART4, MST158,
	MSTP158, Nin one binding protein, NIN1/RPN12 binding protein 1 homolog S. cerevisiae, nob1,
	NOB1_HUMAN, NOB1P, Phosphorylation regulatory protein HP-10, Protein ART-4, Protein ART4,
	PSMD8 binding protein 1, PSMD8BP1, RNA binding protein NOB1, RNA-binding protein NOB1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5007746 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	Background: NOB1P is a 412 amino acid nuclear protein that is involved in proteosome
	biogenesis and is required for the final step in 18S rRNA maturation. NOB1P contains a PIN
	domain, which functions as a nuclease in nonsense-mediated mRNA decay and is required for
	pre-rRNA cleavage. NOB1P interacts with Rent2, which is involved in nonsense-mediated decay
	of mRNAs containing premature stop codons. Expressed in placenta, spleen, endothelial cells,
	liver and lung, NOB1P is essential for the synthesis of 40S ribosome subunits. Supression of the
	gene encoding NOB1P inhibits the processing of the 20S pre-rRNA to the mature 18S rRNA,
	therefore leading to accumulation of high levels of 20S pre-rRNA with degradation
	intermediates.
Gene ID:	28987
Pathways:	Protein targeting to Nucleus
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months