antibodies -online.com





Datasheet for ABIN5007754

anti-OPRL1 antibody (AA 1-47) (Alexa Fluor 680)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | OPRL1 | |
| Binding Specificity: | AA 1-47 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This OPRL1 antibody is conjugated to Alexa Fluor 680 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from mouse Nociceptin receptor | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Predicted Reactivity: | Dog,Cow,Pig,Horse,Guinea Pig | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | OPRL1 | |
|-------------------|--------------------------------------|--|
| Alternative Name: | Nociceptin receptor (OPRL1 Products) | |

Target Details

| Background: | Synonyms: KOR3, ORGC, ORL1, Oprl, XOR1, morc, KOR-3, LC132, MOR-C, Nociceptin receptor, | | |
|---------------------|---|--|--|
| | K3 opiate receptor, Kappa-type 3 opioid receptor, Orphanin FQ receptor, Oprl1, Oor | | |
| | Background: G-protein coupled opioid receptor that functions as receptor for the endogenous | | |
| | neuropeptide nociceptin. Ligand binding causes a conformation change that triggers signaling | | |
| | via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream | | |
| | effectors. Signaling via G proteins mediates inhibition of adenylate cyclase activity and calcium | | |
| | channel activity. Arrestins modulate signaling via G proteins and mediate the activation of | | |
| | alternative signaling pathways that lead to the activation of MAP kinases. Plays a role in | | |
| | modulating nociception and the perception of pain. Plays a role in the regulation of locomotor | | |
| | activity by the neuropeptide nociceptin. | | |
| Gene ID: | 18389 | | |
| UniProt: | P35377 | | |
| Application Details | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |