

## Datasheet for ABIN5007766

## anti-Reticulon 4 antibody (AA 301-400) (AbBy Fluor® 680)



Go to Product page

Overviev	

Overview	
Quantity:	100 μL
Target:	Reticulon 4 (RTN4)
Binding Specificity:	AA 301-400
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Reticulon 4 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence
	(Paraffin-embedded Sections) (IF (p)), Flow Cytometry (FACS)
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from mouse RTN4 protein
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse
Purification:	Purified by Protein A.
Target Details	
Target:	Reticulon 4 (RTN4)

## **Target Details**

<u> </u>	
Background:	Synonyms: ASY, NgA, NOGO, NSP-CL, Nogo-A, Nogo-B, Nogo-C, AA407876, AA409940,
	AA960376, mKIAA0886, mKIAA4153, 1110020G17Rik, C130026I10Rik, Reticulon-4, Neurite
	outgrowth inhibitor, Nogo protein, Rtn4, Kiaa0886
	Background: Developmental neurite growth regulatory factor with a role as a negative regulator
	of axon-axon adhesion and growth, and as a facilitator of neurite branching. Regulates neurite
	fasciculation, branching and extension in the developing nervous system. Involved in down-
	regulation of growth, stabilization of wiring and restriction of plasticity in the adult CNS.
	Regulates the radial migration of cortical neurons via an RTN4R-LINGO1 containing receptor
	complex. May inhibit BACE1 activity and amyloid precursor protein processing.
Gene ID:	68585
UniProt:	Q99P72
Pathways:	Neurotrophin Signaling Pathway, Regulation of Cell Size, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months