



[Go to Product page](#)

Datasheet for ABIN5007809

anti-NPDC1 antibody (AA 75-180) (Alexa Fluor 680)

Overview

Quantity:	100 µL
Target:	NPDC1
Binding Specificity:	AA 75-180
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NPDC1 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NPDC1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat
Purification:	Purified by Protein A.

Target Details

Target:	NPDC1
Alternative Name:	NPDC1 (NPDC1 Products)

Target Details

Background:	<p>Synonyms: AI314472, CAB1, Neural proliferation differentiation and control protein 1, NPDC-1, NPDC1, NPDC1_HUMAN.</p> <p>Background: NPDC-1 (Neural Proliferation Differentiation and Control-1) is expressed in neurons once they have stopped dividing and begun to differentiate. NPDC-1 is transported from the Golgi apparatus via vesicles before becoming internalized by endosomes at the cell membrane. NPDC-1 interacts with Cdk2, D-type cyclins, and the transcription factor E2F1. This interaction can lead to an increased replication time, and might have implications in final neural differentiation and apoptosis. NPDC-1 has been shown to colocalize with synaptic vesicle proteins: synaptophysin, synaptobrevin 2, and Rab3 GEP (Rab3 GTP/GDP exchange protein). One function of NPDC-1 is to regulate retinoic acid-mediated events by directly interacting with retinoid receptors. The amino acid sequence of NPDC-1 is highly conserved between mouse, rat, and human.</p>
-------------	---

Application Details

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months