

Datasheet for ABIN5007940

anti-NDEL1 antibody (AA 241-345) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	NDEL1
Binding Specificity:	AA 241-345
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NDEL1 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NUDEL
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NDEL1
Alternative Name:	NUDEL (NDEL1 Products)

Target Details

Background:	Synonyms: A. nidulans, DKFZp451M0318, ENDOOLIGOPEPTIDASE A, EOPA, MITAP 1, MITAP1,NDEL1, Mitosin associated protein 1, Mitosin associated protein MITAP1, Mitosin-associated protein 1, Ndel 1, Ndel1, NDEL1_HUMAN, Nuclear distribution gene E like homolog 1, Nuclear distribution protein nudE like 1, Nuclear distribution protein nudE-like 1, NUDE like protein, NudE nuclear distribution gene E homolog like 1 A. nidulans, NudE nuclear distribution gene E homolog like 1, NUDEL, Protein Nudel. Background: Nudel is important for normal cortical development. It is involved in microtubule organization, nuclear translocation, and neuronal positioning in concert with various other factors (including Lis1, Pafah1b1, Pafah1b2, dynein, dynorphin A and cdk5). Western blot analysis of mouse tissues shows abundant expression of Nudel in brain and testis, and much lower expression in heart, liver, kidney, and skeletal muscle. In fractionated rat brain, Nudel and Lis1 are both found in fractions enriched for postsynaptic density proteins. Immunostaining of embryonic day 18 mouse brain sections revealed staining of migrating neurons and thalamocortical axons of the intermediate zone of the developing cerebral cortex, as well as several other developing brain regions. The deduced protein contains 345 amino acids and has a calculated molecular mass of 38.4 kDa. It has a coiled coil motif (residues 19 to 201), followed by several potential phosphorylation sites for casein kinase II, protein kinase C or CDK5. Nudel shares about 50 % identity with mouse and human NUDE proteins.
Gene ID:	81565
Pathways:	Regulation of Cell Size

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months