# antibodies -online.com





## anti-NUP107 antibody (AA 501-600) (Alexa Fluor 680)



Go to Product page

| $\sim$ |           |      |    |   |
|--------|-----------|------|----|---|
|        | $ V \cap$ | r\/I | 19 | ٨ |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | NUP107  |
| Binding Specificity: | AA 501-600  |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This NUP107 antibody is conjugated to Alexa Fluor 680 |
| Application:         | Western Blotting (WB), Flow Cytometry (FACS)          |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Nuclear pore complex protein Nup107 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Human   |
| Predicted Reactivity: | Mouse,Rat,Pig,Horse,Rabbit  |
| Purification:         | Purified by Protein A.  |

### Target Details

| Target:           | NUP107   |
|-------------------|--|
| Alternative Name: | NUP107/Nuclear Pore Complex Proteins (NUP107 Products)                                 |
| Background:       | Synonyms: 107 kDa nucleoporin, NPC proteins, NU107_HUMAN, Nuclear pore complex protein |

Nup107, Nucleoporin 107 kDa, Nucleoporin Nup107, NUP 107, NUP 84, NUP107, NUP153, NUP155, NUP84, NUP98.

Background: Nuclear pore complexes (NPCs) are the channels for the bi-directional movement of macromolecules between the nucleus and cytoplasm, and contain more than 100 different subunits. Many of them belong to a family called nucleoporins, which are characterized by the presence of O-linked N-acetylglucosamine moieties and a distinctive pentapeptide repeat (XFXFG). Nuclear pore complex protein Nup107 (Nucleoporin Nup107) is an essential part of the nuclear pore complex which is composed of Nup133, Nup160, Nup107 and Nup96. Nup107 is active in assembling peripheral proteins into this complex. The protein, located on both the nuclear and cytoplasmic sides of the pore, is important in RNA export.

Pathways:

Protein targeting to Nucleus

FCM 1:20-100

#### **Application Details**

Application Notes:

| Application Notes. | FGW 1.20-100   |
|--------------------|--|
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |